Contents of Volume 37

NUMBER 1 JANUARY 1999

Perturbation of ^{15}N chemical shifts following *N*-oxidation of the alkaloid cryptolepinone determined by ^{1}H–^{15}N GHSQC and GHMBC at natural abundance
G. E. MARTIN, C. E. HADDEN, J. R. BLINN, M. H. M. SHARAF, A. N. TACKIE AND P. L. SCHIFF, JR. 1

Technical aspects of an efficient multiple-solvent suppression pulse sequence
C. DALVIT, G. SHAPIRO, J.-M. BÖHLEN AND T. PARELLA 7

Ring inversion of fluorocyclohexand in its solid thiourea inclusion compound
R. K. HARRIS, A. NORDON AND K. D. M. HARRIS 15

Multinuclear magnetic resonance studies under pressure of gases in supercritical media employing a novel titanium-sapphire high pressure cell with pressure sensor
S. GAEMERS, H. LUYTEN, J. M. ERNSTING AND C. J. ELSEVIER 25

Methyl β-substituent effect on NMR ^{17}O chemical shifts in two-coordinated oxygen atoms: DFT GIAO and NBO and experimental studies
J. E. PERALTA, R. H. CONTRERAS, O. E. TAURIAN, F. S. ORTIZ, D. G. DE KOWALEWSKI AND V. J. KOWALEWSKI 31

A new-water-soluble and lipid-insoluble spin probe: application to the study of aqueous sucrose solutions
P. S. BELTON, L. H. SUTCLIFFE, D. G. GILLIES, X. WU AND A. I. SMIRNOV 36

Electronic structure and phosphorus shielding: transition between PN double and triple bonding in iminophosphines RO–P=NMes*
D. GUDAT, E. NIECKE, G. GROSSMANN AND K. KRÜGER 43

Application of selective HSQC experiment to measure interglycosidic heteronuclear long-range coupling constants in cyclodextrins
P. FORGO AND V. T. D'SOUZA 48

Elucidation of the conformations and absolute configurations of enantiomerically pure tetralin derivatives
G. TÓTH, A. SIMON, T. LINKER, F. REBIEN, J. KRAUS AND G. BRINGMANN 53

Complete assignment of the ^{1}H and ^{13}C NMR spectra of the Coccinellidae-defensive alkaloids myrrhine, precoccinelline, hippodamine, their *N*-oxides and the corresponding hydrochlorides
B. LEBRUN, J. C. BRAEKMAN AND D. DALOZE 60

Notes

Synthesis and ^{1}H and ^{13}C NMR study of stereochemistry of (*E*)-2-acetyl-3-aryl-7-arylidene-3,3a,4,5,6,7-hexahydroindazoles
V. VIJAYABASKAR, S. PERUMAL AND M. J. E. HEWLINS 65

***E*/*Z*-isomerism of semicyclic amidinium salts**
S. HARTMANN, V. BRECHT AND A. W. FRAHM 69

^{1}H NMR study of heterocyclic symmetric and assymetric Troger's base analogs containing the acridine ring
M. DEMEUNYNCK, C. FONTAINE AND J. LHOMME 73

Stereoselective hetero Diels–Alder reactions: structure determination of new xanthenedione derivatives by NMR spectroscopy and x-ray chrystallography
A. GIRAUD, P. VANELLE AND L. GIRAUD 77

Reference Data

^{1}H and ^{13}C NMR assignments of conformationally defined 6-s-*cis*-retinoids
D. D. MUCCIO, Y. C. LEE, M. ALAM AND W. J. BROUILLETTE 82

^{1}H and ^{13}C chemical shifts of some tetracyclic and pentacyclic acridinone derivatives
J. P. HANOUN, S. MOREL, V. PIQUE, J. P. GALY AND R. FAURE 86

Papers in Press, 90

NUMBER 2 FEBRUARY 1999

^{13}C chemical shifts in the aryl complexes PdX(p-RC$_6$H$_4$)(TMEDA). Is the PdX(TMEDA)-fragment a substantial π-doner?
M. TSCHOERNER, R. W. KUNZ AND P. S. PREGOSIN 91

Characterization of metabolites in plant extracts of *Ancistrocladus likoko* by HPLC coupled on-line with ^{1}H NMR spectroscopy
G. BRINGMANN, M. RÜCKERT, W. SAEB AND V. MUDOGO 98

NMR vs. molecular modelling II—steroid chemical shift calculations
W. B. SMITH 103

NMR vs. molecular modelling III—a DFT–GIAO study of a series of 2-X-fluorobenzenes
W. B. SMITH 107

NMR vs. molecular modelling IV—conformational analysis of menthone and isomenthone imines
W. B. SMITH AND C. A. AMEZCUA 110

Characterization of a conformationally flexible system: NMR spectroscopic study of 1,5-diphenyl-2,4-pentanedione
J. T. PULKKINEN, R. LAATIKAINEN, J. J. VEPSÄLÄINEN AND M. J. AHLGRÉN 119

Determination by 1D NOE studies of the stereochemistry of a new tricyclic 2-hydroxytetrahydrofuran derivative obtained by hydroformulation of (1*R*)-(–)-myrtenol
S. SIROL, J.-P. GORRICHON, P. KALCK, P. M. NIETO AND G. COMMENGES 127

Synthesis and multinuclear NMR study of (*E*)-s-*trans*-1-acetyl-5-aryl-3-styryl-2-pyrazolines
V. VIJAYABASKAR, S. PERUMAL, S. SELVARAJ, A. LYCKA, R. MURUGAN AND M. BALASUBRAMANIAN 133

Notes

Complete ^{1}H and ^{13}C NMR assignments of diosgenyl saponins
X. W. HAN, H. YU, X. M. LIU, X. BAO, B. YU, C. LI AND Y. Z. HUI 140

Conformational analyses of model structures for intramolecular glycoside bond formation
A. GEYER, U. HUCHEL AND R. R. SCHMIDT 145

Synthesis and spectral study of *N*-2-(3′-phthalidyl)phthalazin-1-one
Y. CHEN, W. HUA, M. LÜ AND J. PAN 149

Structure determination and detailed ^{1}H, ^{13}C NMR assignments of triterpenes from *Mimusops elengi*
N. P. SAHU, K. KOIKE, Z. JIA, N. B. MONDAL, B. ACHARI AND T. NIKAIDO 152

Reference Data

^{13}C NMR study of *N*-acyl-*S*-homoserine lactone derivatives
W. LAO, S. KJELLEBERG, N. KUMAR, R. DENYS, R. W. READ AND P. STEINBERG 157

Complete assignment of the ^{1}H, ^{13}C and ^{19}F NMR Spectra of fluoro-substituted [3*H*]-naphthopyranes
S. DELBAERE, Y. TERAL, C. BOCHU, M. CAMPREDON AND G. VERMEERSCH 159

^{13}C NMR spectral assignments of 3α,3′α-bis(arylcarboxy)-5β-cholan-24-oic acid 1,2-ethanediol diesters: new lithocholic acid-based molecular clefts
J. TAMMINEN, K. LAPPALAINEN, K. LAIHIA, P. MÄNTTÄRI, H. SALO AND E. KOLEHMAINEN 163

Papers in Press, 164

NUMBER 3 MARCH 1999

DFT–GIAO and DFT–NBO studies of the origin of ^{19}F NMR shielding effects in alkyl fluorides
W. ADCOCK, D. LÜNSMANN, J. E. PERALTA AND R. H. CONTRERAS 167

Selenium-77 and carbon-13 high-resolution solid-state NMR studies of selenomethionine
M. J. POTRZEBOWSKI, R. KATARZYŃSKI AND W. CIESIELSKI 173

^{1}H and ^{13}C NMR characterization of poly[3-(6-methoxyhexyl)-2,2′-bithiophene]
D. IAROSSI, A. MUCCI, L. SCHENETTI, P. COSTA BIZZARRI, C. DELLA CASA AND M. LANZI 182

NMR study of methyl 3,4,6-tri-*O*-acetyl-2-deoxy-2-[alkylureido]-β-D-glucopyranosides
I. WAWER, M. WEYCHERT, J. KLIMKIEWICZ, B. PIEKARSKA-BARTOSZEWICZ AND A. TEMERIUSZ 189

^{15}N NMR study of isoquinoline alkaloids
R. MAREK, O. HUMPA, J. DOSTÁL, J. SLAVÍK AND V. SKLENÁŘ 195

Measurement of the deuterium relaxation times in double-labeled (^{13}C/^{2}H) thymidine and 2′-deoxyadenosine and in the selectively labeled DNA duplex: $^{5'}$d(^{1}C^{2}G^{3}A^{4}T^{5}T^{6}A^{7}A^{8}T^{9}C^{10}G)$_{2}^{3'}$
T. V. MALTSEVA, A. FÖLDESI AND J. CHATTOPADHYAYA 203

Influence of γ-hydrogens (hydrogens three bonds away) in calculating ^{13}C chemical shifts of acyclic alkanes and alcohols
A. SAIKIA AND I. SURYANARAYANA 214

Stereochemical arrangement of spiro[4.5]decane derivatives by ^{15}N and ^{17}O NMR
A. ARIZA-CASTOLO AND J. GODOY-REYES 219

Orientation of electric field gradient and chemical shift tensors in $BiVO_4$ at room temperature by ^{51}V NMR
P. R. BODART, Y. DUMAZY J. P. AMOUREUX AND C. FERNANDEZ 223

Intramolecular electric field effect on a ^{1}J(CH) NMR spin–spin coupling constant. An experimental and theoretical study
D. G. DE KOWALEWSKI, V. J. KOWALEWSKI, J. E. PERALTA, G. ESKUCHE, R. H. CONTRERAS, A. L. ESTEBAN, M. P. GALACHE AND E. DÍEZ 227

Carbon-13 and nitrogen-15 solution- and solid-state NMR of some sulphonated azobenzene dyestuffs
A. M. CHIPPENDALE, G. McGEORGE, R. K. HARRIS AND C. M. BRENNAN 232

Notes

Complete ^{1}H and ^{13}C NMR Spectra Assignment of 6,7,8,9-tetrahydro-7,7-dimethyl-dioxolo[5,4-*b*]acridin-9-one using FUCOUP
C. J. NASCIMENTO, E. R. CRUZ AND J. FIGUEROA-VILLAR 239

Deuterium-induced carbon-13 isotope shifts in nitrogen-containing benzoheterocyclic systems
M. S. MORALES-RÍOS, H. CERVANTES-CUEVAS, I. SALGADO-ESCOBAR AND P. JOSEPH-NATHAN 243

Book reviews, 246; Announcement, 248; Papers in press, 249

NUMBER 4 APRIL 1999

New, stable β-phosphorus-labelled pyrrolidine nitroxides from nitrones for magnetometry: An ESR investigation
D. L. HAIRE, E. G. JANZEN, G. CHEN, V. J. ROBINSON AND I. HRVOIC 251

NMR of enaminones. Part 6—^{17}O and ^{13}C NMR study of tautomerization in Schiff bases
J.-C. ZHUO 259

NMR spectroscopic diffusion, chemical shift and linewidth measurements of low-affinity binding of ibuprofen enantiomers to human serum albumin
Y. MA, M. LIU, X.-A. MAO, J. K. NICHOLSON AND J. C. LINDON 269

Proton transfer equilibrium in Schiff bases derived from 5-nitrosalicyladehyde. A study of deuterium isotope effects on ^{13}C NMR chemical shifts
Z. ROZWADOWSKI AND T. DZIEMBOWSKA 274

Interaction of [Ile7]angiotensin III with dodecylphosphocholine micelles: NMR and molecular dynamics studies
A. B. PATEL, S. SRIVASTAVA AND R. S. PHADKE 279

^{1}H and ^{13}C NMR study of the paramagnetic chiral macrocyclic lanthanide complexes
J. LISOWSKI 287

Complete assignments of ^{1}H and ^{13}C NMR spectra of the pentacyclic triterpene hederagenin from *Nigella sativa* Linn
B. S. JOSHI, K. L. SINGH AND R. ROY 295

^{14}N NMR single crystal study of hexamethylenetetramine
Y.-S. KYE AND G. S. HARBISON 299

^{13}C chemical shift tensors and secondary structure of poly-L-alanine by solid state two-dimensional spin-echo NMR and *ab initio* chemical shielding calculation
N. ASAKAWA, M. TAKENOIRI, D. SATO, M. SAKURAI AND Y. INOUE 303

Notes

Complete assignment of the ^{13}C NMR spectra of a series of 5,8-disubstituted 4,4-dimethyl-anthracene-1,9,10(4*H*)-triones
R. ARAYA-MATURANA, B. K. CASSELS, T. DELGADO-CASTRO, C. HURTADO-GUZMÁN AND C. JULLIAN 312

Internal rotation processes in *endo-cis-N*-(*o*-tolyl)bicyclo[2.2.1]heptene-2,3-dicarboximide and its oxidation products
M. P. A. CAMPOS, L. BERGTER, C. H. T. DE PAULA E SILVA AND P. R. SEIDL 317

^{13}C NMR spectra of triangular μ-oxoruthenium acetate clusters revisited: HETCOR study of two pyridine and 4,4′–bipyridine derivatives
H. E. TOMA, A. D. P. ALEXIOU, S. NIKOLAOU AND S. DOVIDAUSKAS 322

Effects of *N*-oxidation on the ^{15}N chemical shifts in the *Strychnos* alkaloids strychnine and brucine
C. E. HADDEN, B. D. KALAZNY, R. H. ROBINS AND G. E. MARTIN 325

Reference Data

^{1}H and ^{13}C NMR chemical shift assignment of some ^{3}*H*-napthto[2,1-*b*]pyrans
R. DEMADRILLE, C. MOUSTROU, M. CAMPREDON, G. GIUSTI AND R. FAURE 328

Announcement, 331; Papers in Press, 332

NUMBER 5 MAY 1999

^{13}C, ^{19}F and ^{77}Se NMR study of vicinal (*E*)-fluoro(organylseleno)olefins and [(*E*)-fluoroalkenyl]di-organylselenonium salts
H. POLESCHNER, M. HEYDENREICH AND R. RADEGLIA 333

Investigations on a flexible prolyl-endopeptidase inhibitor in solution by NMR techniques
B. PODÁNYI, S. BOKOTEY, K. KÁNAI, M. FEHÉR AND I. HERMECZ 346

Single-crystal EPR study of the thermally accessible $^{3}B_u$ state of *trans*-$(CNSSS)_2^{2+}$
A. BERCES, G. D. ENRIGHT, G. E. McLAURIN, J. R. MORTON, K. F. PRESTON, J. PASSMORE AND D. J. WOOD 353

NMR spectroscopy in environmental chemistry: ^{1}H and ^{13}C NMR parameters of tricyclic poly-chlorinated C_{10} hydrocarbons and their oxy derivatives based on two-dimensional NMR techniques
E. KOLEHMAINEN, J. KOIVISTO, K. LAIHIA, R. KAUPPINEN AND J. PAASIVIRTA 359

Structural characterisation and dynamic NMR studies of a new peracylated macrocyclic diterpene
G. GÜNTHER, T. MARTINEK, G. DOMBI, J. HOHMANN AND A. VASAS 365

Determination of the relative configuration of the C-19 to C-29 fragment of amycin B by ^{1}H and ^{13}C NMR investigations
F. HOYER, G. HABERMEHL, H. DUDDECK AND G. TÓTH 371

Note

Solid-state formation of centrosymmetric cage dimeric 4-aryl-1,4-dihydropyridines via non-symmetric *syn*-dimers studied by ^{13}C CPMAS NMR spectroscopy
A. HILGEROTH, G. HEMPEL, U. BAUMEISTER AND D. REICHERT 376

Rapid Communications

Selective 1D $^{1}H \rightarrow ^{13}C \rightarrow ^{13}C$ polarization transfer experiments for small organic molecules
L. POPPE, S. GRÖGER AND J. ZABLOCKI 382

Low level long-range ^{1}H–^{15}N heteronuclear shift correlation at natural abundance using a combination of 2D and selective 1D NMR methods with submicro NMR probe technology
C. E. HADDEN AND G. E. MARTIN 385

Book reviews, 389; Papers in press, 391

NUMBER 6 JUNE 1999

Application of LC–NMR to the identification of bulk drug impurities in GART inhibitor AG2034
B. C. M. POTTS, K. F. ALBIZATI, M. O'NEIL JOHNSON AND J. PROSSER JAMES 393

Conformational analysis of crown ether analogs in solution: *cis*-cyclohexano-8-crown-3 as studied via low-temperature ^{13}C and ^{1}H NMR spectroscopy and molecular mechanics calculations
G. W. BUCHANAN, A. B. DRIEGA, R. C. LAISTER AND K. BOURQUE 401

Multivariate correlation between ^{13}C and ^{31}P NMR spectral data on dissolved lignin and the combustion properties of kraft black liquor
P. MALKAVAARA, R. ALÉN AND E. KOLEHMAINEN 407

Determination of the sign of long-range ^{1}H–^{13}C scalar coupling constants to non-protonated carbons: application to assignment
Y. KUMAKI AND K. HIKICHI 413

Tetrakis(trimethylstannyl)germanium, the first observation of indirect nuclear ^{119}Sn–^{73}Ge spin–spin coupling
B. WRACKMEYER AND P. BERNATOWICZ 418

GIAO–DFT prediction of accurate NMR parameters in selected glucose derivatives
T. KUPKA, G. PASTERNA, P. LODOWSKI AND W. SZEJA 421

Structure of silylated benzohydroxamic acids
J. SCHRAML, M. KVÍČALOVÁ, L. SOUKUPOVÁ AND V. BLECHTA 427

Notes

Complete ^{1}H and ^{13}C NMR assignments of stephalic acid
E. BURGUEÑO-TAPIA, L. R. HERNÁNDEZ, P. JOSEPH-NATHAN AND M. SALMÓN 430

Chemical transformations on a weakly acidic catalyst studied by ^{15}N NMR: butanone oxime on silicalite-1
W. O'NEIL PARKER, JR 433

Predominance of inductive over resonance substituent effect on ^{33}S NMR chemical shifts of 4-substituted phenyl-4′-methylphenacyl sulfones
R. GAWINECKI, E. KOLEHMAINEN, A. ZAKRZEWSKI, K. LAIHIA, B. OŚMIAŁOWSKI AND R. KAUPPINEN 437

Reinvestigation of the substituent effects of fluorine on the ^{1}H NMR chemical shifts of the adjacent methylene group in fluorinated [2.2]paracyclophanes
L. ERNST AND K. IBROM 441

Reference Data

^{1}H and ^{13}C NMR study of 6-substituted 2-naphthyl methyl sulphides
S. PERUMAL, S. SELVARAJ AND M. J. E. HEWLINS 445

^{1}H and ^{13}C NMR spectroscopy analysis of acetonides of D-glucitol derivatives
L. KATO AND R. M. BRAGA 447

Papers in Press, 450

NUMBER 7 JULY 1999

Review Article

A practical approach to the implementation of selectivity in homonuclear multidimensional NMR with frequency selective-filtering techniques. Application to the chemical structure elucidation of complex oligosaccharides
C. ROUMESTAND, C. DELAY, J. A. GAVIN AND D. CANET 451

Conformational analysis of the biphenyl moiety of dimer liquid crystals in the solid state by high-resolution solid-state ^{13}C NMR and *ab initio* GIAO-CHF calculations
R. N. SHIMIZU, H. KUROSU, I. ANDO, A. ABE AND H. FURUYA 479

ESR study of the copper(II)–glycylglycine equilibrium system in fluid aqueous solution. Computer analysis of overlapping multispecies spectra
T. SZABÓ-PLÁNKA, A. ROCKENBAUER AND L. KORECZ 484

^{1}H, ^{13}C and ^{15}N NMR study of some triazolo-, and tetrazolo pyridazines and thioxotriazolopyridines
P. CMOCH, L. STEFANIAK, E. MELZER, S. BAŁONIAK AND G. A. WEBB 493

Stereochemical effects in some acronycine derivatives
E. MIKROS, S.MITAKU, A.-L. SKALTSOUNIS, F. LIBOT, F. TILLEQUIN AND M. KOCH 498

Review Article

A practical approach to the implementation of selectivity in homonuclear multidimensional NMR with frequency selective-filtering techniques. Application to the chemical structure elucidation of complex oligosaccharides
C. ROUMESTAND, C. DELAY, J. A. GAVIN AND D. CANET 451

Conformational analysis of the biphenyl moiety of dimer liquid crystals in the solid state by high-resolution solid-state ^{13}C NMR and *ab initio* GIAO-CHF calculations
R. N. SHIMIZU, H. KUROSU, I. ANDO, A. ABE AND H. FURUYA 479

ESR study of the copper(II)–glycylglycine equilibrium system in fluid aqueous solution. Computer analysis of overlapping multispecies spectra
T. SZABÓ-PLÁNKA, A. ROCKENBAUER AND L. KORECZ 484

^{1}H, ^{13}C and ^{15}N NMR study of some triazolo-, and tetrazolo pyridazines and thioxotriazolopyridines
P. CMOCH, L. STEFANIAK, E. MELZER, S. BAŁONIAK AND G. A. WEBB 493

Stereochemical effects in some acronycine derivatives
E. MIKROS, S.MITAKU, A.-L. SKALTSOUNIS, F. LIBOT, F. TILLEQUIN AND M. KOCH 498

Note

Complete assignment of ^{1}H and ^{13}C NMR data and establishment of the relative stereochemistry of C-1-functionalized 2,4-dimethyl-8-oxabicyclo[3.2.1]oct-6-en-3-one derivatives
A. M. MONTAÑA, P. M. GRIMA AND F. GARCÍA 507

Rapid Communication

Application of a high-resolution superconducting NMR probe in natural product structure determination
T. M. LOGAN, N. MURALI, G. WANG AND C. JOLIVET 512

Papers in Press, 516

NUMBER 8 AUGUST 1999

ACCORD-HMBC: advantages and disadvantages of static versus accordion excitation
G. E. MARTIN, C. E. HADDEN, R. C. CROUCH AND V. V. KRISHNAMURTHY 517

Rapid acquisition of ^{1}H–^{13}C and ^{1}H–^{15}N heteronuclear chemical shift correlation data at the sub-milligram level using SMIDG (submicro inverse-detection gradient) NMR probe technology
G. E. MARTIN, C. E. HADDEN, A. N. TACKIE, M. M. H. SHARAF AND P. L. SCHIFF, JR 529

An unusual temperature dependence of the EPR parameters of the copper(II) and oxovanadium(IV) bis(acetylacetonate) complexes
B. D. HOWES, C. KÜHLMEYER, R. POGNI AND R. BASOSI 538

Indirect nuclear spin–spin coupling in tetrachlorotetraborane(4) and halogenated polyhedral phospha- and arsaboranes
W. KELLER, W. HAUBOLD AND B. WRACKMEYER 545

***Ab initio* calculation of NMR shielding in phosphaalkenes X—P=CY_2**
A. B. ROZHENKO, W. W. SCHOELLER AND M. I. POVOLOTSKII 551

Three-dimensional solution structures of two DNA dodecamers through full relaxation matrix analysis
I. BERTINI, A. CLEMENTE, I. ROMBECK, A. ROSATO, P. TURANO, B. LIPPERT AND F. QUADRIFOGLIO 564

^{59}Co second-order quadrupolar effects in the ^{13}C cross-polarization magic angle spinning NMR spectra of the cobaltocenium salts $[Cp^*_2Co]^+[PF_6]^-$ and $[Cp^*CpCo]^+[PF_6]^-$
H. HEISE, F. H. KÖHLER, E. B. BROUWER, R. K. HARRIS AND S. STEUERNAGEL 573

Notes

Influence of the introduction of fluorine atoms on the ^{13}C NMR spectra of acrylic esters
L. MARIVAL-HODEBAR, M. TORDEUX AND C. WAKSELMAN 579

Principal axis orientation of the ^{17}O hyperfine tensor of NO_2
R. G. HAYES 583

Conformational flexibility of disaccharides investigated by multiple field carbon-13 nuclear spin relaxation
P. SÖDERMAN AND G. WIDMALM 586

Reference Data

^{1}H and ^{13}C chemical shifts of 4,5-disubstituted acridine derivatives
S. ISSMAILI, V. PIQUE, J.-P. GALY AND R. FAURE 591

Total assignment of ^{1}H and ^{13}C NMR spectra of pregnenolone and progesterone haptens using 2D NMR spectroscopy
S. S. KORDE, R. KATOCH, R. A. UDASI AND G. K. TRIVEDI 594

Assignment of the ^{13}C NMR signals of some 8-azaadenine, 2,6-diaminopurine and 2,6-diamino-8-azapurine carbonucleosides
M. TEIJEIRA, C. TERÁN AND E. URIARTE 598

^{13}C NMR relaxation study of monoaminopyridines in D_2O and $CDCl_3$–DMF solutions
F. CESARE MARINCOLA, E. CADONI, G. GELLI, A. LAI AND P. BELTRAME 600

^{1}H and ^{15}N chemical-shift assignments of a carboxy-terminal functional domain of the bacteriophage P22 scaffolding protein
Y. SUN AND N. RAMA KRISHNA 602

Determination of the structure of 3-acylmethylene-2,3-dihydrofurans with the aid of 2D NOESY spectra: ^{1}H and ^{13}C NMR chemical shifts assignment
J. G. SOŚNICKI AND Z. WICHERT-TUR 605

Papers in Press, 608

NUMBER 9 SEPTEMBER 1999

NMR detection of oriented association via dilution shifts in tetramethylsilane solvent. 1. *n*-Alkyl halides
G. V. D. TIERS 609

NMR detection of oriented association via dilution shifts in tetramethylsilane solvent. 2. Aliphatic esters
G. V. D. TIERS, J. STEVENS AND W. L. STEBBINGS 613

Multi-dimensional NMR structural analyses of purified lipid X and lipid A (endotoxin)
A. A. RIBEIRO, Z. ZHOU AND C. R. H. RAETZ 620

Novel selective 1D experiments based on heteronuclear *J* cross-polarization
T. PARELLA, J. BELLOC, F. SÁNCHEZ-FERRANDO AND A. VIRGILI 631

Combined use of complete lineshape analysis of 1D spectra subjected to reference deconvolution and linear prediction, 2D-EXSY spectra and a double fitting method for the study of chemical exchange. Application to an eight-site exchange system
P. S. DENKOVA AND V. S. DIMITROV 637

Gradient-enhanced proton–fluorine NOE experiments
J. T. GERIG 647

Elucidation of the stereochemistry of octahydroisoquinoline derivatives by NMR
I. GHIVIRIGA, P. Q. BOTTARI AND T. HUDLICKY 653

Characterization of biguanides by ^{15}N NMR spectroscopy
B. CLEMENT AND U. GIRRESER 662

Extremely large one-bond $^{31}P^{13}C$ and two-bond $^{31}P^{18}O$ isotope shifts in $[(H_3C)_2CH]_3Si{-}O{-}C{\equiv}P$, a new representative of λ^3-phosphaalkynes, and isotope shifts of several derivatives and some λ^3-phosphaalkenes
G. HECKMANN, G. BECKER AND H. KRAFT 667

Notes

NMR characterization of 1,1-selenocystine with 92% ^{77}Se enrichment
M. SALZMANN, E. M. STOCKING, L. A. SILKS, III AND H. SENN 672

Two known bis-indole alkaloids isolated from *Tabernaemontana laeta*: complete ^1H and ^{13}C chemical shift assignments
W. L. B. MEDEIROS, I. J. C. VIEIRA, L. MATHIAS, R. BRAZ-FILHO, K. Z. LEAL, E. RODRIGUES-FILHO AND J. SCHRIPSEMA 676

Reference Data

^1H and ^{13}C NMR spectra for a series of arylmaleamic acids, arylmaleimides, arylsuccinamic acids and arylsuccinimides
J. TRUJILLO-FERRARA, R. SANTILLAN, H. I. BELTRÁN, N. FARFÁN AND H. HÖPFL 682

NMR spectral data of acyclovir prodrugs
H. GAO AND A. K. MITRA 687

Book reviews, 690; Papers in press, 691

NUMBER 10 OCTOBER 1999

Temperature dependence study of chlorine NQR frequency and spin–lattice relaxation time in 2-amino-3,5-dichloropyridine
MOHAMMED HOSSEIN A. A., K. P. RAMESH, N. RAJENDRA KUMAR AND J. RAMAKRISHNA 693

Coordination equilibrium—a clue for fast water exchange on potential MRI contrast agents?
E. TÓTH, O. M. NI DHUBHGHAILL, G. BESSON, L. HELM AND A. E. MERBACH 701

Analysis of cross-polarisation dynamics between ^1H and ^{19}F in Viton fluoroelastomer using solid-state ^{19}F magic angle spinning and $^1H{\rightarrow}^{19}F$ cross polarization magic angle spinning NMR
S. ANDO, R. K. HARRIS, G. A. MONTI AND S. A. REINSBERG 709

Comparison of 1.7 mm submicro and 3 mm micro gradient NMR probes for the acquisition of ^1H–^{13}C and ^1H–^{15}N heteronuclear shift correlation data
G. E. MARTIN AND C. E. HADDEN 721

ESR measurements of the partitioning of some new spin probes in *n*-octanol–water
S. E. BOTTLE, D. G. GILLIES, A. S. MICALLEF, D. A. REID AND L. H. SUTCLIFFE 730

Spin delocalisation in radical anions containing two phosphaalkene groups: an EPR study
A. AL BADRI, A. JOUAITI AND M. GEOFFROY 735

NMR spectroscopy in environmental chemistry: ^{1}H and ^{13}C NMR chemical shift assignments of chlorinated dibenzothiophenes based on two dimensional NMR techniques and *ab initio* MO and DFT/GIAO calculations
E. KOLEHMAINEN, J. KOIVISTO, V. NIKIFOROV, M. PERÄKYLÄ, K. TUPPURAINEN, K. LAIHIA, R. KAUPPINEN, S. A. MILTSOV AND V. S. KARAVAN 743

Notes

Distinguishing diastereomeric organic compounds based on the concerted use of ^{3}J(C,H) and ^{3}J(N,H) values
S. GRÖGER, J. A. JOSEY AND L. POPPE 748

Neoadifolin, a new indole alkaloid from the heartwood of *Adina cordifolia*
B. BALÁZS, G. TÓTH, P. S. RAO, T. NARENDER, N. SUBBA RAO, GY. HORVÁTH AND H. DUDDECK 751

Two cardenolides from *Calotropis procera*
A. G. HANNA, M. H. A. ELGAMAL, N. A. M. MORSY, H. DUDDECK, J. KOVÁCS AND G. TÓTH 754

NMR study of tautomerism of benzofuroxan and its sulphur and selenium analogues
P. CMOCH AND W. SCHILF 758

Rapid Communication

Application of a high-resolution superconducting NMR probe in natural product structure determination†
T. M. LOGAN, N. MURALI, G. WANG AND C. JOLIVET 762

† This paper, original reference *Magn. Reson. Chem.* **37,** 512 (1999), is republished here following technical problems with the printing of the original article.

NUMBER 11 NOVEMBER 1999

ESR of new β-phosphorus nitroxides from aldo- and keto-nitrones
D. L. HAIRE, E. G. JANZEN AND I. HRVOIC 767

EPR and MO calculation studies on α-aminoanthraquinone anion radicals in aprotic solvents
V. VATANEN, J. M. ELORANTA AND M. VUOLLE 774

^{1}H and ^{13}C NMR study of quaternary benzo[*c*]phenanthridine alkaloids
R. MAREK, J. TOUŠEK, J. DOSTÁL, J. SLAVÍK, R. DOMMISSE AND V. SKLENÁŘ 781

Etoposide: conformational and hydration features
K. ZHANG, A. FRANZ, G. C. HILL, C. M. McCALLUM AND M. J. MINCH 788

Multiple field carbon-13 NMR relaxation of calix[4]arene in solution
M. SHULTES, R. EISENHAUER, J. J. DECHTER, M. JOHANSSON, P. KUMAR AND J. KOWALEWSKI 799

Recognition of chirality in nitroxides using EPR and ENDOR spectroscopy
P. SCHULER, F.-M. SCHABER, H. B. STEGMANN AND E. JANZEN 805

Conformational analysis of 3′-*S*-PO_3-linked ribo- and deoxyribodinucleoside monophosphates
A. P. G. BEEVERS, E. M. WITCH, B. C. N. M. JONES, R. COSSTICK, J. R. P. ARNOLD AND J. FISHER 814

An improved *J*-multiplied method for the measurement of $^{3}J(H^{N},H^{\alpha})$ from the two-dimensional ^{15}N–^{1}H correlation spectrum
P. PERMI, I. KILPELÄINEN AND S. HEIKKINEN 821

Solvent effects in the structure dereplication of caffeoyl quinic acids
G. F. PAULI, U. KUCZKOWIAK AND A. NAHRSTEDT 827

Stuctures of two novel isomeric triterpenoid saponins from *Anthocephalus cadamba*
N. P. SAHU, K. KOIKE, Z. JIA, B. ACHARI, S. BANERJEE AND T. NIKAIDO 837

Differential broadening caused by dipolar interaction with ^{1}H in ^{13}C MAS NMR signals split by residual dipolar interaction with ^{14}N
S. HAYASHI 843

Notes

^{17}O NMR chemical shifts: a simple and useful rule for substituent additivity on the oxygen atoms with a coordination number of two
M. A. P. MARTINS, N. ZANATTA, H. G. BONACORSO, G. M. SIQUEIRA AND A. F. C. FLORES **852**

NMR assignments of ellagic acid derivatives
X.-C. LI, H. N. ELSOHLY, C. D. HUFFORD AND A. M. CLARK **856**

Complete assignment of the ^{1}H and ^{13}C NMR spectra of secoisolariciresinol diglucoside, a mammalian lignan precursor isolated from *Linum usitatissimum*
S. CHIMICHI, M. BAMBAGIOTTI-ALBERTI, S. A. CORAN, V. GIANNELLINI AND B. BIDDAU **860**

Book reviews, 864; Papers in press, 866

NUMBER 12 DECEMBER 1999

Structure determination of unusual mixed-ligand complexes $[L_{OMe}InTp]X$ ($X = [InCl_4]^-$, $[PF_6]^-$) in solution by 2D ^{1}H NOESY NMR spectroscopy
W. KLÄUI, N. LIEDTKE AND W. PETERS **867**

Application of EPR spectroscopy to the characterization of magnetic interactions in thermally decomposed coal
B. PILAWA, A. B. WIĘCKOWSKI AND M. LEWANDOWSKI **871**

Electron density distribution in 2-nitro-5-methylimidazole derivatives studied by NMR–NQR double resonance
J. N. LATOSIŃSKA, J. SELIGER AND B. NOGAJ **878**

Studies of the electronic structure of 4-*N*-cytosine derivatives by NMR–NQR double resonance spectroscopy
J. N. LATOSIŃSKA, J. SELIGER, V. S. GRECHISHKIN AND J. SPYCHALA **881**

Configurational and conformational NMR study of enantiopure 2,2-dimethyl-1-(1-naphthyl)propanol via its carbamate derivatives
M. POMARES, X. GRABULEDA, C. JAIME, A. VIRGILI, A. ÁLVAREZ-LARENA AND J. F. PINIELLA **885**

Application of 2D TOCSY to determination of the stoichiometry of strand associations in multi-stranded DNA structures
P. K. PATEL AND R. V. HOSUR **891**

NMR spectra of nitrogen-containing compounds. Correlations between experimental and GIAO calculated data
A. DOKALIK, H. KALCHHAUSER, W. MIKENDA AND G. SCHWENG **895**

Notes

Protonation sites on unsymmetrical 1,3-diamines: a nuclear magnetic resonance study of 3-aminotropanes
A. H. LEWIN, J. P. BURGESS, G. SUN AND L. D. FUDALA **903**

Total assignment of ^{1}H and ^{13}C NMR spectra of tectol and tecomaquinone I
T. L. G. LEMOS, S. M. O. COSTA, O. D. L. PESSOA AND R. BRAZ-FILHO **908**

Reference Data

^{13}C, ^{31}P and ^{1}H NMR of some dioxaphosphinopyridines and pyridoazaphosphinines
A. FERREIRA, M. BIGAN AND D. BLONDEAU **912**

^{13}C NMR and ^{31}P NMR spectral assignment of new β-phosphonylated hydrazones
A. BEN AKACHA, S. BARKALLAH AND H. ZANTOUR **916**

Author index, 921; Subject index, 924; Papers in press, 928; Volume contents, iii

SPECIAL ISSUE DECEMBER 1999

Guest Editors' Foreword
Magnetic resonance and zeolites
J. KÄRGER AND D. MICHEL i

Editorial
Magnetic resonance of porous materials
H. GÜNTHER ii

Review Articles
^{129}Xe NMR overview of xenon physisorbed in porous solids
M.-A. SPRINGUEL-HUET, J.-L. BONARDET, A. GÉDÉON AND J. FRAISSARD S1

Applications of laser-polarized ^{129}Xe under continuous flow
E. BRUNNER S14

Research Articles
ESR observation of redox processes on benzene-loaded NaPdY, NaPtY and NaCeY zeolites
G. HUBNER AND E. RODUNER S23

^{29}Si NMR study of base-catalyzed polymerization of dimethyldiethoxysilane
S. E. RANKIN AND A. V. MCCORMICK S27

Time-resolved ^{13}C MAS NMR investigation of chemical reactions: the MTG process
T. MILDNER, H. ERNST, D. FREUDE, J. KÄRGER AND U. WINKLER S38

Studies of Fe(III) incorporated into $AlPO_4$-20 by X- and W-band EPR spectroscopies
D. ARIELI, D. E. W. VAUGHAN, K. G. STROHMAIER, H. THOMANN, M. BERNARDO AND D. GOLDFARB S43

Characterization of offretite during hydrothermal treatment by high-resolution solid-state ^{29}Si magic angle spinning NMR and ^{27}Al triple-quantum magic angle spinning NMR
P. LENTZ, A. P. CARVALHO, L. DELEVOYE, C. FERNANDEZ, J.-P. AMOUREUX AND J. B.NAGY S55

Multinuclear solid-state NMR study of cadmium- and lead-exchanged LTA zeolites
A. ELDEWIK, J. M. HOOK, N. K. SINGH AND R. F. HOWE S63

Applications of ^{27}Al multiple quantum magic angle spinning NMR to aluminophosphate molecular sieves
P. R. BODART, J. P. AMOUREUX, M. PRUSKI, A. BAILLY AND C. FERNANDEZ S69

Characterization of basic guest compounds on solid catalysts by ^{13}C CP/MAS NMR spectroscopy of surface methoxy groups
U. SCHENK, M. HUNGER AND J. WEITKAMP S75

Combined quasi-elastic neutron scattering and NMR study of hydrogen diffusion in zeolites
N.-K. BÄR, H. ERNST, H. JOBIC AND J. KÄRGER S79

Diffusion behavior of the benzene molecule in faujasite-type zeolite studied by double quantum filtered NMR
Y.-H. CHEN AND L.-P. HWANG S84

EPR study of NO adsorption–desorption behaviour on Lewis acid sites in NaA zeolites
T. RUDOLF, A. PÖPPL, W. BRUNNER AND D. MICHEL S93

A set of two-dimensional solid-state NMR experiments providing topological information for structure refinement of TMP-GaPO
V. MUNCH, F. TAULELLE, T. LOISEAU, G. FÉREY, A. K. CHEETHAM, S. WEIGEL AND G. D. STUCKY S100

***n*-Pentane hopping in zeolite ZK-5 studied with ^{13}C NMR**
P. C. M. M. MAGUSIN, D. SCHURING, E. M. VAN OERS, J. W. DE HAAN AND R. A. VAN SANTEN S108

Characterization of Brønsted and Lewis acidity in zeolites by solid-state NMR and the recent progress in the REDOR technique
A. L. BLUMENFELD AND J. J. FRIPIAT S118

^{1}H NMR studies of the mobility of olefins adsorbed in zeolites of type NaX
W. BÖHLMANN, D. MICHEL AND J. ROLAND S126

Terminal and bridging methoxyls on zeolites detected by ^{13}C MAS NMR
V. BOSÁČEK, R. KLIK, F. GENONI, G. SPANO, F. RIVETTI AND F. FIGUERAS S135

NMR study of adsorption techniques for organic compounds on silica surfaces
H. GÜNTHER, S. OEPEN, M. EBENER AND V. FRANCKE S142

^{1}H NMR relaxation measurements in highly concentrated water protein solutions
R. OLECHNOWICZ, W. MASIERAK, J. BODURKA AND A. GUTSZE S147

Temperature dependence of the EPR linewidth of ultramarine blue
A. B. WIĘCKOWSKI, W. WOJTOWICZ AND J. ŚLIWA-NIEŚCIÓR **S150**

Historical Perspective

A short history of nuclear magnetic resonance spectroscopy and of its early years in Germany
H. PFEIFER **S154**

Papers in Press, S160